

Engraved by T. Sansom

William Curtis,

FLORA LONDINENSIS, BOTANICAL MAGAZINE &c.



Engraved by T. Sansom

William Curtis,

FLORA LONDINENSIS, BOTANICAL MAGAZINE &c.

2. 3. B.k

#### THE

### BOTANICAL MAGAZINE;

OR,

### Flower-Garden Displayed:

#### IN WHICH

The most ornamental FOREIGN PLANTS, cultivated in the Open Ground, the Green-House, and the Stove, will be accurately represented in their natural Colours.

#### TO WHICH WILL BE ADDED,

Their Names, Class, Order, Generic and Specific Characters, according to the celebrated Linn Eus; their Places of Growth, and Times of Flowering:

TOGETHER WITH

THE MOST APPROVED METHODS OF CULTURE.

#### A WORK

#### Intended for the Use of

Such LADIES, GENTLEMEN, and GARDENERS, as wish to become fcientifically acquainted with the Plants they cultivate.

#### LONDON:

Printed by FRY and COUCHMAN, Upper-Moorfields,
For W. CURTIS, at his BOTANIC GARDEN, Lambeth-Marsh;
And fold by all Bookfellers, Stationers, and News-Carriers,
in Town and Country.

M DCC LXXXVII.

MVSEVM BRITANNICVM

BOTANICAL MINCARINES

Flower Gurden Diplayed:



### KAKAKAKAKAKAKAKAKAKA

### PREFACE.

THE present periodical publication owes its commencement to the repeated solicitations of several Ladies and Gentlemen, Subscribers to the Author's BOTANIC GARDEN, who were frequently lamenting the want of a work, which might enable them, not only to acquire a systematic knowledge of the Foreign Plants growing in their gardens, but which might at the same time afford them the best information respecting their culture—in sact, a work, in which Botany and Gardening (so far as relates to the culture of ornamental Plants) or the labours of Linnæus and Miller, mitchappily be combined.

In compriance with their wishes, he has endeavoured to present them with the united information of both authors, and to illustrate each by a set of new figures, drawn always from the living plant, and coloured as near to nature, as the impersection of colouring will admit.

He does not mean, however, to confine himself solely to the Plants contained in the highly esteemed works of those luminaries of Botany and Gardening, but


#### PREFACE.

but shall occasionally introduce new ones, as they may slower in his own garden, or those of the curious in any part of Great-Britain.

At the commencement of this publication, he had no defign of entering on the province of the Florist, by giving figures of double or improved Flowers, which sometimes owe their origin to culture, more frequently to the sportings of nature; but the earnest entreaties of many of his Subscribers, have induced him so far to deviate from his original intention, as to promise them one, at least, of the Flowers most esteemed by Florists.

The encouragement given to this work, great beyond the Author's warmest expectations, demands his most grateful acknowledgements, and will excite him to persevere in his humble endeavours to render Botany a lasting source of rational amusement, and public utility.

BOTANIC GARDEN, Lambeth-Marsh,





Published at the direct to H. Burnis Brennis Berton Lambeth March with

### [ 1 ]

#### IRIS PERSICA. PERSIAN IRIS.

\*\*\*\*\*\*\*

Class and Order.

#### TRIANDRIA MONOGYNIA.

#### Generic Character.

Corolla 6-partita: Petalis alternis, reflexis. Stigmata petaliformia.

Specific Character and Synonyms.

IRIS Persica corolla imberbi, petalis interioribus brevissimis patentissimis. Lin. Syst. Vegetab. p. 79. Sp. Pl. p. 59.

IRIS bulbofa præcox minus odora Perfica variegata. Morif. bift. 2. p. 357.


XIPHIUM Perficum. Miller Diet. ed. 6. 410.

The Persian bulbous Flower-de-luce. Parkins. Parad. p. 172.

A native of Persia. Flowers in February and March. Its beauty, early appearance, and fragrant blossoms, make it highly esteemed by all lovers of slowers; like the Hyacinth or Narcissus it will blow within doors in a water-glass, but stronger in a small pot of sand, or sandy loam; a few slowers will scent a whole apartment: it will also blossom in the open air, but requires warmth and shelter; it is propagated by offsets and seeds; the best slowering roots are imported from Holland, they bear forcing well; and hence this plant may be had to slower a full month or six weeks in succession.

. Parkinson remarks, that in his time (1629) it was very rare, and seldom bore slowers.

tajal salek 196 abisaltajal おというの のではないかのからないか The state of the s The state of the s A Charles Company of the least d Parks I'm





### RUDBECKIA PURPUREA. PURPLE RUDBECKIA.



Class and Order.

SYNGENESIA POLYGAMIA FRUSTRANEA.

Generic Character.


Receptaculum paleaceum, conicum. Pappus margine quadridentato. Calyx duplici ordine fquamarum.


Specifie Character and Synonyms,


RUDBECKIA purpurea foliis lanceolato-ovatis alternis indivifis, radii petalis bifidis. Lin. Syst. Vegetab. p. 651. Sp. Pl. p. 1280.

DRACUNCULUS virginianus latifolius, petalis florum longissimis purpurascentibus. Moris. bist. 3. p. 42. s. 6. t. 9. f. 1.

This species differs from the other plants of the genus, in the colour of its outermost petals, which are long, narrow, purple, and pendulous, and not unaptly resemble small pieces of red tape. Notwithstanding it is a native of the warm climates Carolina and Virginia, it succeeds very well with us in an open border; but, as Mr. Miller very justly observes, it will always be prudent to shelter two or three plants under a common hot-bed frame in winter, to preserve the kind, because in very severe winters, those in the open air are sometimes killed. It slowers in July. As it rarely ripens its seeds with us, the only mode of propagating it, is by parting the roots; but in that way the plant does not admit of much increase.







Problem in the Lines by It have nothing combin Lambeth March, 1766.

### HELLEBORUS HYEMALIS. WINTER HELLEBORE OF ACONITE.

\*\*\*\*\*\*\*\*\*\*\*

Class and Order.
POLYANDRIA POLYGYNIA\*.

Generic Character.

Calyx o. Petala 5 five plura. Nectaria bilabiata, tubulata. Capfulæ polyspermæ erectiusculæ.

Specific Character and Synonyms.


HELLEBORUS byemalis flore folio infidente. Lin. Syst. Vegetab. p. 431. Sp. Pl. p. 783.


ACONITUM unifolium bulbosum. Banb. Pin. 183. The Winter's Wolfesbane. Park. Parad. p. 214.

Grows wild in Lombardy, Italy, and Austria, affects mountainous situations, slowers with us in February, and hence is liable to be cut off by severe frosts. "Is propagated by offsets, "which the roots send out in plenty. These roots may be taken up and transplanted any time after their leaves decay, which is generally by the beginning of June till October, when they will begin to put out new fibres; but as the roots are small and nearly the colour of the ground, so if care is not taken to search for them, many of the roots will be lest in the ground. These roots should be planted in small clusters, otherwise they will not make a good appearance, for single slowers scattered about the borders of these small kinds are scarce seen at a distance; but when these and the Snowdrops are alternately planted in bunches, they will have a good effect, as they slower at the same time, and are much of a size." Miller's Gard. Dist.

<sup>\*</sup> Most of the Hellebores vary greatly in the number of their pistils, which in general are too few to justify the placing those plants in the order Polygynia.

Henry will dries in susping H CONTRACTA TO ASSESSED TO This world the buildy in as demicionary by the property 





Tublish I as the Lit directs by W. Corn. Bestonic Garden Lambeth March 1986.

#### [4]

## CYCLAMEN COUM. ROUND-LEAV'D CYCLAMEN.

\*\*\*\*\*

Class and Order.

PENTANDRIA MONOGYNIA.

Generic Character.

Corolla rotata, reflexa, tubo brevissimo fauce prominente. Bacca tecta capsula.

Specific Character and Synonyms.

CYCLAMEN Coum foliis orbiculatis planis, pediculis brevibus, floribus minoribus. Miller's Dia.

CYCLAMEN hyemale orbiculatis foliis inferius rubentibus purpurascente flore; Coum Herbariorum. Hort. reg. Paris. Herm. Cat.

CYCLAMEN orbiculato folio inferne purpurascente. Baub. Pin. p. 307.


The common round-leav'd Sowebread. Park. Parad. p. 198.


Grows wild in many parts of Italy and Germany, and is fometimes found with white flowers; if the feason be mild, or the plants sheltered from the inclemency of the weather, this species will flower as early as February, or much earlier by artificial heat.

As it grows naturally in woods and shady places, it will thrive best in a mixture of bog-earth and loam placed in a north border; if planted in the open border, it will require to be covered with a hand-glass during winter, and in the spring, when in bloom; the more usual method with gardeners is to preserve them in pots in a common hot-bed frame, the advantage of this method is that they may, at any time, be removed to decorate the parlour or the study.

The plants of this genus admit of but little increase by their roots; the best method of propagating them is by seed, which should be sown soon after they are ripe in boxes or pots, and covered about half inch deep, placing them where they may have only the morning-sun, till the beginning of September, when they may be removed to a warmer exposure.

1 1 Creating Comma Round Carly p MANA MOVO come closely would and taged out a price promise necessition and trader interest for the second of the second and produced by the second of the second production and a second government as early the change or making caller by have seemen nationally in wheels and thedy places, it will with breath and bud three you land which a he field with premiers that it is the story of the property for the story of the story 





# ERYTHRONIUM DENS CANIS. DOGS-TOOTH, or DOGS-TOOTH VLOLET.

\*\*\*\*\*\*

Class and Order.
HEXANDRIA MONOGYNIA.

Generic Character.

Corolla 6-petala, campanulata: Nectario tuberculis 2-petalorum alternorum basi adnatis.

Specific Character and Synonyms.

ERYTHRONIUM Dens Canis. Lin. Syst. Vegetab. p. 269. Sp. Pl. p. 437.

Dens Canis latiore rotundioreque folio. Baub. Pin. 87.

Dogs-Tooth with a pale purple flower. Park. Parad. p. 194.

Of this genus Mr. Miller makes too species; Linnæus, perhaps with more propriety, only one, for breadth of leaves or colour of flowers can scarcely be considered as sufficient to constitute a specific difference.


It is found in the gardens with purple flowers of two different tints, also with white and yellow blossoms, grows naturally in Hungary and some parts of Italy, and blows in the open border


at the beginning of April.

5

"They are propagated by offsets from their roots. They love a shady situation and a gentle loamy soil, but should not be too often removed. They may be transplanted any time after the beginning of June, when their leaves will be quite decayed, till the middle of September; but the roots should not be kept very long out of the ground, for if they shrink it will often cause them to rot. The roots of these flowers should not be planted scattering in the borders of the slower-garden, but in patches near each other, where they will make a good appearance." Miller's Gard. Dist.

... O'sza C k spy čateryki orrange the second section of the affirmation (through the second control of the second seco small of water has since the alle water Table danger 1 Allered to see wiged with the the state of het agent an fall t man and the second second second





the link is a the Act direct by Wilson's Botonic Carden Lambeth March 1786

#### NARCISSUS MINOR. LEAST DAFFODIL.



Class and Order.

HEXANDRIA MONOGYNIA.

Generic Character.

Petala 6, æqualia: Nectario infundibuliformi, 1-phyllo. Stamina intra nectarium.

Specific Character and Synonyms.

NARCISSUS minor spatha unislora, nectario obconico erecto crispo sexsido æquante petala lanceolata. Lin. Sp. Pl. p. 415. Syst. Vegetab. p. 262.

NARCISSUS parvus totus luteus. Baubin. Pin. 53.

The least Spanish yellow bastard Dasfodil. Park. Parad. p. 105.


We are not a little surprised that Mr. Miller should have taken no notice of the present species, as it must have been in the English gardens long before his time, being mentioned by Parkinson in his Garden of pleasant Flowers: it is nearly related to the Pseudo-Narcissus, but differs from it in many particulars except size, vid. Lin. Sp. Pl. and Parkinson above quoted.


Though its bloffoms are not so large as those of the other species, yet when the roots are planted in a cluster, they make a very pretty shew, and have this advantage, that they slower somewhat earlier than any of the others.

Like the common Daffodil it propagates very fast by the roots, and will thrive in almost any soil or situation.

Though a native of Spain, it is feldom injured by the feverity of our climate.


Marriage anno." Learn Dersonal HOLDER TO SOURCE STRUCK COME etion school in 19th and a selection of the 19th of 19th and additional string products of the decision Control resultant parties of Texas 10 Miles Control Special State of the St eyed bloods with a law self ball on the way as a law way The most winder to the least to the being dentioned by the state of th seem representation of the second representation and the second A STATE OF THE PARTY OF THE PAR





Published as the Art directs by Wilartis Botanic Garden Lambeth March 1787

## CYNOGLOSSUM OMPHALODES. BLUE NAVELWORT.



Class and Order.

PENTANDRIA MONOGYNIA.

Generic Character.

Corolla infundibuliformis, fauce claufa fornicibus. Semina depressa interiore tantum latere stylo assixa.

Specific Character and Synonyms.

CYNOGLOSSUM Omphalodes repens, foliis radicalibus cordatis\*, Lin. Sp. Pl. p. 193. Syft. Vegetab. p. 157. Scopoli Fl. Carn. p. 124. t. 3.


SYMPHYTUM minus borraginis facie. Baub. Pin. 259.

BORAGO minor verna repens, folio lævi. Moris. bist. 3. p. 437. s. 11. t. 26. sig. 3.

A native of Spain, Portugal, and Carniola, and an inhabitant of woods and shady situations, slowers in March and April: in the autumn it puts forth trailing shoots, which take root at the joints, whereby the plant is most plentifully propagated; thrives best under a wall in a North border.

<sup>&</sup>quot; Stolones repunt non caulis florifer, cui folia ovalia, et minime cordata. "TOURNEFORTIUS feparavit a Symphito, et dixit Omphallodem pumilam vernam, symphyti folio, fed bene monet LINNÆUS folam fructus afperitatem aut glabritiem, non sufficere ad novum genus construendum." Scopoli Fl. Carn. p. 124.

Olivers of Chamberra i de d'avent de la constitue 





Published as the Act directs by W. Garais, Bosonis Garden Lambeth Marsh 1986

# HELLEBORUS NIGER. BLACK HELLEBORE, or CHRISTMAS ROSE.

\*\*\*\*\*\*\*\*\*

Class and Order.

POLYANDRIA POLYGYNIA.

Generic Character.

Calyx nullus. Petala 5 five plura. Nectaria bilabiata, tubulata. Capfulæ polyfpermæ, erectiufculæ.

Specific Character and Synonyms.

HELLEBORUS niger scapo sub-bisloro subnudo, foliis pedatis. Lin. Syst. Vegetab. p. 431. Sp. Pl. p. 783.

HELLEBORUS niger flore roseo, Baub. Pin. 186.

The true Black Hellebore, or Christmas slower. Parkins. Parad. p. 344.

As our Publication feems likely to fall into the hands of fuch as are totally unacquainted with Botany, or botanical writings, it must plead as an apology for our often explaining many circumstances relative to plants, which may be well known to

adepts in the science.

This plant derives its first name from the black colour of its roots, its second from its early slowering, and the colour of its petals, which though generally milk-white on their first appearance, yet have frequently a tint of red in them, which increases with the age of the blossom and finally changes to green; in some species of Hellebore, particularly the viridis, the slower is green from first to last.

Black Hellebore grows wild on the Appenine and other

mountains, preferring fuch as are rocky.


If the weather be unufually mild, it will flower in our gardens, in the open border, as early as December and January; it may indeed be confidered as the herald of approaching spring.

Like most other alpine plants, it loves a pure air, a situation moderately most, and a soil unmanured: as the beauty of its slowers is apt to be destroyed by severe frosts, it should be covered during the winter with a hand-glass, or if it be treated in the manner recommended for the round-leav'd Cyclamen, it may be had to slower in still greater perfection.

It is propagated by parting its roots in autumn: neither this

species nor the byemalis thrive very near London.

HEREROLDE MOCE. BLACK HELERORE. Later stellar 1 which is the piece of the could be belower, to be later -shop with a burney congression of the tell to the tel Attract the field one of the flower flower. Table that to alone on year of an extra property of the end o ni burets du la company de la manufactory from the best of the best of the best of the eith a the season to be promined to be promined to be promined this the state of the state of the state of the state of · number





Published as the Act directs by W. Ourtis Novanis Carden Lambeth Marsheyer.

## [ 9 ]

### IRIS PUMILA. DWARF IRIS.



Class and Order.

#### TRIANDRIA MONOGYNIA.

Generic Character.

Corolla fex-partita: Petalis alternis, reflexis. Stigmata petali-

Specific Character and Synonyms.

IRIS pumila corollis barbatis, caule foliis breviore unissoro.

Lin. Syst. Vegetab. p. 78. Sp. Plant. p. 56. Jacq. Fl.

Austr. t. 1.

CHAM ÆIRIS minor flore purpureo. Baub. Pin. 33.

The leffer purple dwarf Flower-de-luce. Park. Parad. p. 186.

Gardeners, in former days, not having that profusion of plants to attend to and cultivate, which we can at present boast, appear to have been more solicitous in increasing generally the varieties of the several species; accordingly, we find in the Paradisus terrestris of the venerable Parkinson, no less than six varieties of this plant\*, most of which are now strangers to the Nursery Gardens. We may observe, that varieties in general not being so strong as the original plant, are consequently much sooner lost.

The Iris pumila grows wild in many parts of Hungary, affects open and hilly fituations, and flowers in our gardens in the month of April; it is a very hardy plant, and will thrive in almost any soil or fituation; is propagated by parting its roots in autumn.

_	The lesser purple dwarf Flower-de-luce with white blossoms,							
		-		-	ftraw colour ditto.			
			-	_	pale blue ditto.			
		-		-	blush-coloured ditto.			
i				-	yellow variable ditto.			
		_		_	blue variable ditto, and			

the purple dwa


LESS FOR LESS AND LES


Cardeners, in former days, to this age that producion of oldness to attend to and countwing, while, we constitute posterior book to be not more softs were in recreasing generalise the serieties of the accept dispersal of pectodary of one of the serieties of the accept dispersal of the serieties of the plant the serieties which are now indices to be varieties of the plant to be serieties of the serieties of the

The shis probab sport within mone parts of Hought yell stalls open and belt fruncises a read discourt in all graduating the month of the part of the section of the section

er had blackers	 sistematical in	ingwi - on	100 MT
ATAT COLOUR LAND.			
	 40000000	-	
a fluor hadden objection of	 		-
the all the world			
and bld in the relie			

the purple dwarf Sea Flower-de less of the come nuther, is reduced to





Published as the Act directs by W. Guras, Rotanic Garden Lambath March 1767

## ANEMONE HEPATICA. HEPATICA, or Noble Liverwort.

\*\*\*\*\*\*\*

Class and Order.

POLYANDRIA POLYGYNIA.

Generic Character.

Calyx nullus. Petala 6. g. Semina plura.

Specific Character and Synonyms.

ANEMONE Hepatica foliis trilobis integerrimis. Lin. Syft. Vegetab. p. 424. Sp. Pl. p. 758. Fl. Suec. n. 480.

TRIFOLIUM hepaticum flore simplici et pleno. Bauh. Pin. 339. Red Hepatica or noble Liverwort. Park. Parad. p. 226.

Dillenius, Miller, and some other authors, make a distinct genus of the *Hepatica*: Linnæus unites it with the *Anemone*, observing, that though it differs from the *Anemone* in having a calyx, yet that calyx is at some distance from the flower, and partakes more of the Nature of an Involucrum, which is not uncommon to the Anemonies.


The Hepaticas, as Parkinfon observes, flower foon after the winter Hellebore, " and making their pride appear in winter,

" are the more welcome early guests."

It is found wild in its fingle state, with red, blue, and white slowers, in the woods and shady mountains of Sweden, Germany, and Italy; the red variety with double slowers is the one most commonly cultivated in our gardens; the double blue is also not unfrequent; the single white is less common; and the double white Miller never saw, yet admits that it may exist spontaneously, or be produced from seed: Parkinson mentions a white variety with red threads or stamina.

According to Miller, this plant delights in a loamy foil, and in an eastern position where it may have only the morning fun: the single forts are easily raised from seed; the double, increased by parting the roots, which ought to be done in March when they are in bloom; they should not be divided into very small heads: these plants, if often removed and parted, are apt to die, but lest undisturbed for many years, they will thrive exceedingly, and become very large roots.

CRAME - ALLOY STREET RESTRICTION 





Published as the Act directs by W Curtes Hotanic Garden, Lambeth March 1767

### ERICA HERBACEA. HERBACEOUS HEATH.

\*\*\*\*\*\*\*\*

Class and Order.

OCTANDRIA MONOGYNIA.

Generic Character.

Calyx 4-phyllus. Corolla 4-fida. Filamenta receptaculo inferta. Antheræ bifidæ. Capfula 4-locularis.

Specific Character and Synonyms.

ERICA berbacea antheris muticis exfertis, corollis oblongis, ftylo exferto, foliis quaternis, floribus fecundis, Lin. Syst Vegetab. p. 306. carnea Sp. Pl. ed. 3. p. 504.

ERICA carnea. Jacq. Fl. Austr. v. 1. tab. 32

ERICA procumbens herbacea. Baub. Pin. p. 486.

Since the days of Mr. Miller, who, with all his imperfections, has contributed more to the advancement of practical gardening than any individual whatever, our gardens, but more especially our green-houses, have received some of their highest ornaments from the introduction of a great number of most beautiful Heaths: the present plant, though a native of the Alps and mountainous parts of Germany, is of modern introduction here, what renders it particularly acceptable, is its hardiness and early flowering; its blossoms are formed in the autumn, continue of a pale green colour during the winter, and expand in the spring, slowering as early as March, especially if kept in a green-house, or in a common hot-bed frame, which is the more usual practice.

It may be propagated by feeds or cuttings, the latter is the most ready way of increasing this and most of the other species of the genus: when the cuttings have struck root, they should be planted in a mixture of fresh loam and bog earth, either in


the open border, under a wall, or in pots.

The name of berbacea, which Linnæus has given to this plant, is not very characteristic, but it should be observed, that Linnæus in this, as in many other instances, has only adopted the name of some older botanist; and it should also be remembered, that in genera, where the species are very numerous, it is no easy matter to give names to all of them that shall be perfectly expressive.

This species does not appear to us to be specifically different

from the mediterranea.

AND ASSESSED FOR STATE OF THE SAME ASSESSED. Indiana and the second terms to the Agreement state of SA Paint State of the Control of the STORES OF THE PARTY. 





y h

n

I Somethy del at fouly.

Published as the set directs by M. Curris Bottonic Gordon Lamboth March

## DODECATHEON MEADIA. MEAD'S DODECA-THEON, OF AMERICAN COWSLIP.

\*\*\*\*\*\*

Class and Order.

PENTANDRIA MONOGYNIA.

Generic Character.

Corolla rotata, reflexa. Stamina tubo infidentia. Capfula unilocularis, oblonga.

Specific Character and Synonyms.

DODECATHEON Meadia. Lin. Syst. Vegetab. p. 163. Sp. Plant. p. 163.

MEADIA Catesb. Car. 3. p. 1. t. 1. Trew. Ebret. t. 12.

AURICULA ursi virginiana floribus boraginis instar rostratis, cyclaminum more reslexis. Pluk. alm. 62. 1. 79. f. 6.

This plant grows spontaneously in Virginia and other parts of North America, from whence, as Miller informs us, it was sent y Mr. Banister to Dr. Compton, Lord Bishop of London, in those curious garden he first saw it growing in the year 1709.

It is figured by Mr. Catesby, in his Natural History of Caroina, among the natural productions of that country, who belowed on it the name of *Meadia*, in honour of the late Dr. Mead, a name which Linnæus has not thought proper to dopt as a generic, though he has as a trivial one.

" It flowers the beginning of May, and the feeds ripen in July, foon after which the stalks and leaves decay, so that the

roots remain inactive till the following spring.

"It is propagated by offsets, which the roots put out freely when they are in a loofe moist soil and a shady situation; the best time to remove the roots, and take away the offsets, is in August, after the leaves and stalks are decayed, that they may be fixed well in their new situation before the frost comes on. It may also be propagated by seeds, which the "plants"

" plants generally produce in plenty; these should be fown " in autumn, foon after they are ripe, either in a shady moist " border, or in pots, which should be placed in the shade; in " the fpring, the plants will come up, and must then be kept clean from weeds; and, if the feafon proves dry, they must be frequently refreshed with water: nor should they be exposed to the sun; for while the plants are young, they are very impatient of heat, fo that I have known great numbers " of them destroyed in two or three days, which were growing " to the full fun. These young plants should not be transplanted till the leaves are decayed, then they may be care-" fully taken up and planted in a fhady border, where the foil " is loofe and moist, at about eight inches distance from each other, which will be room enough for them to grow one " year, by which time they will be strong enough to produce " flowers, fo may then be transplanted into some shady borders " in the flower-garden, where they will appear very ornamental " during the continuance of their flowers." Miller's Gard. Dia.

Conditional that a resident of the first that the

THE REPORT OF THE PARTY OF THE PARTY OF

BADIA Tatel. Co. a real and

entors the state of in pt Xre Small dayle, with any market beginning to ng to be with a mater motor court, and the said per tarbullet, describe an aire per fig. in a epersonal transfer of the second secon oil enterminate entre al company option de la company de la co ch ne ce of a part which is to prife the major through ers tal a.



Published as the Act directs by W.Carnis Botanic Garden Lambeth March 1787.

Sounded

# CORONILLA GLAUCA. SEA-GREEN, OF DAY-SMELLING CORONILLA.

\*\*\*\*\*\*

Class and Order.

#### DIADELPHIA DECANDRIA.

Generic Character.

Calyx bilabiatus: 3: dentibus superioribus connatis.

Vexillum vix alis longius. Legumen isthmis interceptum.

Specific Character and Synonyms.

CORONILLA glauca fruticosa, foliolis septenis, obtusissimis, stipulis lanceolatis. Lin. Syst. Vegetab. p. 557. Sp. Pl. 1047.

CORONILLA maritima, glauco folio. Tournef. inft. 650.


COLUTEA scorpioides maritima, glauco folio. Baub. Pin. 397. prodr. 157.

This charming shrub, which is almost perpetually in blossom, and admirably adapted for nosegays, is a native of the south of France, and a constant ornament to our green-houses.


Linnæus has observed, that the flowers, which in the day time are remarkably fragrant, in the night are almost without scent.

"It is propagated by fowing the feeds in the fpring, either upon a gentle hot-bed, or on a warm border of light earth: when the plants are come up about two inches high, they fhould be transplanted either into pots, or into a bed of fresh earth, at about four or five inches distance every way, where they may remain until they have obtained strength enough to plant out for good, which should be either in pots filled with good fresh earth, or in a warm situated border, in which, if the winter is not too severe, they will abide very well, provided they are in a dry soil." Miller's Gard. Dist.





Nº4



Inthiship as the Act directs by W. Cartis Botanic Cus den Lambeth Mench 1767.

## PRIMULA VILLOSA. MOUNTAIN PRIMULA.

\*\*\*\*\*\*\*\*\*\*\*\*

Class and Order.

#### PENTANDRIA MONOGYNIA.

Generic Character.

Involucrum umbellulæ. Corollæ tubus cylindricus: ore patulo.

Specific Character and Synonyms.

PRIMULA villosa foliis obovatis dentatis villosis, scapo brevissimo multisloro.

PRIMULA villosa. Jacqin Fl. Austr. app. t. 27.


Mr. Miller, in the Sixth Edition of the Abridgment of his Gardener's Dictionary, mentions only four Primulas, exclusive of the Auricula, the two first of which are named erroneously, and of the two last not a syllable is said either as to their place of growth or culture.


The plant here figured, has been introduced pretty generally into the Nursery-Gardens in the neighbourhood of London within these few years: Mr. Salisbury informs me, that a variety of this plant with white flowers, brought originally from the Alps of Switzerland, has for many years been cultivated in a gerden in York him.


cultivated in a garden in Yorkshire.

It is not noticed by Linnæus: Professor Jacquin, in his Flora Austriaca, has figured and described a Primula, which, though not agreeing so minutely as could be wished with the one we have figured, is nevertheless considered by some of the first Botanists in this country as the same species; he gives it the name of villosa, which we adopt, though with us it is so slightly villous as scarcely to deserve that epithet.

It varies in the brilliancy of its colours, flowers in April, and will fucceed with the method of culture recommended for the Round-Leaved Cyclamen.







BARAL L. Warris Rotonic Garden Lambeth March 267.

## [ 15 ] .

## NARCISSUS JONQUILLA. COMMON JONQUIL.



Class and Order.

HEXANDRIA MONOGYNIA.

Generic Character.

Petala fex. Nettario infundibuliformi, monophyllo.

Stamina intra nectarium.

Specific Character and Synonyms.


NARCISSUS Jonquilla spatha multissora, nectario hemisphærico crenato, breviore petalis, soliis semiteretibus. Lin. Spec. Pl. p. 417.

The fragrant Jonquil is a native of Spain, flowers in the open ground, about the latter-end of April, or beginning of May, and will thrive in almost any soil or situation, but prefers, as most bulbs do, a fresh loamy earth; indeed such a soil is savourable to the growth of most plants, as being exempt from a variety of subterraneous insects, which are apt to insest ground which has been long cultivated.

It is found in the gardens with double flowers.

Our plant accords exactly with the description of Linnæus, above quoted, but must be carefully distinguished from some others very similar to it.

the transfer of the sector, and some public for the feet of the discussion and the second





Poblish d to M. Gerrie Botonic Carlos Lumbert March 1987

## [ 16 ]

### IRIS VARIEGATA. VARIEGATED IRIS.

Class and Order.

TRIANDRIA MONOGYNIA.

Generic Character.

Corolla o partita: Petalis alternis, reflexis. Stigmata petali-

Specific Character and Synonyms.

IRIS variegata corollis barbatis, caule subsolioso longitudine foliorum multisloro. Lin. Spec. Pl. p. 56.


IRIS latifolia pannonica, colore multiplici. Baub. Pin. 31.

The yellow variable Flower-de-luce. Parkinson Parad. p. 182.

This species of Iris, inferior to sew in point of beauty, is a native of the hilly pastures of Hungary, and slowers in our gardens in the month of May, and beginning of June. It is a hardy perennial, requires no paricular treatment, and may be easily propagated by parting its roots in Autumn.

Market and the second second second second





#### [ 17 ]

#### CACTUS FLAGELLIFORMIS. CREEPING CEREUS.

\*\*\*\*\*\*\*

Class and Order.

ICOSANDRIA MONOGÍNIA.

Generic Character.

Calyx 1-phyllus, fuperus, imbricatus. Corolla multiplex. Bacca 1-locularis, polyfperma.

Specific Character.

CACTUS flagelliformis repens decemangularis. Lin. Syft. Vegetab. ed. 14. p. 460.


CEREUS flagelliformis. Miller's Gard. Diet. ed. 6. 4to.


Grows spontaneously in South-America, and the West-Indies, slowers in our dry stoves early in June, is tolerably hardy, and will thrive even in a common green-house, that has a flue to keep out the severe frosts.


It is superior to all its congeners in the brilliancy of its colour, nor are its blossoms so sugacious as many of the other species.

No plant is more easily propagated by cuttings; these Miller recommends to be laid by in a dry place for a fortnight, or three weeks, then to be planted in pots, filled with a mixture of loam and lime rubbish, having some stones laid in the bottom of the pot to drain off the moisture, and afterwards plunged into a gentle hot-bed of Tanners bark, to facilitate their rooting, giving them once a week a gentle watering: this business to be done the beginning of July.

It is feldom that this plant perfects its feeds in this country; Miller relates that it has borne fruit in Chelsea gardens.







Times be delationly.

Publish d & Weintis Botanic Garden Lambon Marsh 197.

#### GERANIUM REICHARDI. DWARF GERANIUM.

Class and Order.

MONADELPHIA DECANDRIA.

Generic Character.

Monogyna. Stigmata 5. Fructus irostratus, 5-coccus.

Specific Character and Synonyms.

GERANIUM Reichardi scapis unissoris, sloribus pentandris, foliis subrentsormibus inciso-crenatis.

GERANIUM Reichardi scapis unissoris, foliis plerisque oblongis trilobis vel quinquelobis inciso-crenatis. Lin. Syst. Vegetab. ed. Murr. 14. p 618.


This species of Geranium, so strikingly different from all others at present cultivated in our gardens, has been known for several years to the Nursery-men in the neighbourhood of London, by the name of acaule, a name we should gladly have retained, had not Prosessor Murray described it in the 14th edition of Linnæus's Systema Vegetabilium, under the name of Reichardi, a name he was disposed to give it, in compliment to a French gentleman, who first discovered it in the island of Minorca, and introduced it into the gardens of France.


Linnæus describes many of the Geraniums, as having only five antheræ, though several of those he thus describes have to our certain knowledge ten, the five lowermost of which shedding their pollen first, often drop off, and leave the silaments apparently barren: but in this species (with us at least) there never are more than five, but betwixt each stamen, there is a broad pointed barren silament or squamula, scarcely to be distinguished by the naked eye.

The usual and best practice is to make a green-house plant of this species, though it has been known to remain in the

open ground, during a mild winter, unhurt.

It continues to have a fuccession of blossoms during the greatest part of the summer, and may be propagated either by seed or parting its roots.







#### [ 19 ]

#### HEMEROCALLIS FLAVA. YELLOW DAY-LILY.

\*\*\*\*\*\*\*

Class and Order.

HEXANDRIA MONOGYNIA.

Generic Character.

Corolla campanulata, tubus cylindraceus.

Stamina declinata.

Specific Character and Synonyms.

HEMEROCALLIS flava foliis lineari-subulatis carinatis, corollis slavis. Lin. Syst. Vegetab. ed. 14. p. 339.

LILIUM luteum, asphodeli radice. Baub. Pin. 80.


The Yellow Day-Lily. Parkinf. Parad. p. 148.

This Genus has been called *Hemerocallis*, in English, *Day-Lily*, from the short duration of its blossoms, but these are not quite so fugacious in this species as in the fulva.

It very rarely happens that Linnæus, in his specific character of a plant, has recourse to colour, he has however in this instance; but this seems to arise from his considering them rather as varieties, than species. To us they appear to be perfectly distinct, and in addition to several other characters, the flava is distinguished by the fragrance of its blossoms.

This species is an inhabitant of Hungary and Siberia, and consequently bears our climate exceedingly well; it requires a moist soil, and a situation somewhat shady, and is easily propagated by parting its roots in autumn.

No. 1 College Service 





Publish d by M. Carai. Bottonic Garden, Lambeth March 1767.

J. Sowerby del. et faule.

## GERANIUM PELTATUM. IVY-LEAVED GE-

\*\*\*\*\*\*

Class and Order.

#### MONADELPHIA DECANDRIA.

Generic Character.

Monogyna. Stigmata quinque. Fructus rostratus. 5-coccus.

Specific Character.


GERANIUM peltatum calycibus monophyllis, foliis quinquelobis integerrimis glabris subpeltatis, caule fruticoso. Lin. Syst. Vegetab. ed. 14. p. 613.


GERANIUM africanum, foliis inferioribus afari, superioribus staphidisagriæ maculatis splendentibus et acetosæ sapore. Comm. Præl. 52. t. 2.

A native of Africa, as are most of our shewy Geraniums, is not so tender as many others, and may be propagated very readily from cuttings.

A leaf, having its foot-stalk inserted into the disk or middle part of it, or near it, is called by Linnæus, peltatum, hence the Latin trivial name of this plant. It may be observed, however, that some of the leaves have this character more perfectly than others.

The African Geraniums differ much from the European, in the irregularity of their Petals, but exhibit the character of the Class Monadelphia much better than any of our English ones, having their filaments manifestly united into one body; this species has only 7 filaments bearing antheræ, but 3 barren ones may be discovered upon a careful examination, which makes it of the order Decandria.







I Senato deletjado

Published by W Curic Botanic Garden Lambeth Marsh

#### [ 21 ]

#### IRIS VERSICOLOR. PARTICOLOURED IRIS.



Class and Order.

TRIANDRIA MONOGYNIA.

Generic Character.

Corolla 6-petala, inæqualis, petalis alternis geniculatopatentibus. Stigmata petaliformia, cucullato-bilabiata. Conf. Thunh. Dif. de Iride.


#### Specific Character and Synonyms.


- IRIS versicolor imberbis foliis ensisormibus, scapo tereti flexuoso, germinibus subtrigonis. Lin. Syst. Vegetab. ed. 14. Murr. p. 90. Sp. Plant. ed. 3. p. 57.
- IRIS Americana versicolor stylo crenato. Dill. Elth. 188.
  1. 155. f. 188.


A native of Virginia, Maryland, and Pensylvania, has a perennial root, is hardy, and will thrive in almost any soil or situation; may be increased by parting its roots in autumn.

Our plant is the pilla of Miller, and the versicolor of Miller is, we believe, the sibirica of Linnæus.

This species has, for the most part, a stalk unusually crooked or elbowed, by which it is particularly distinguished. It slowers in June, as do most of this beautiful tribe.







### NIGELLA DAMASCENA. GARDEN FENNEL-FLOWER, LOVE IN A MIST, DEVIL IN A BUSH.

\*\*\*\*\*\*\*\*\*\*

Class and Order.

POLYANDRIA PENTAGYNIA.

Generic Character.

Cal. nullus. Petala 5. Nectaria 5. trifida, intra corollam. Capfulæ 5 connexæ.

Specific Character and Synonyms.

NIGELLA damascena floribus involucro folioso cinctis. Lin.

Syst. Vegetab. ed. 14. Murr. p. 506. Sp. Pl. p. 753.

NIGELLA angustifolia, slore majore simplici cæruleo. Baub.

Pin. 145.


The great Spanish Nigella. Park. Parad. p. 287.


Is an annual, and grows wild among the corn in the fouthern parts of Europe; varies with white and blue flowers, both fingle and double.

"May be propagated by fowing their feeds upon a bed of light earth, where they are to remain (for they feldom fucceed well if transplanted); therefore, in order to have them intermixed among other annual flowers in the borders of the Flower Garden, the feeds should be sown in patches at proper distances: and when the plants come up, they must be thinned where they grow too close, leaving but three or four of them in each patch, observing also to keep them clear from weeds, which is all the culture they require. In July they will produce their flowers, and their feeds will

"ripen in August.
"The season for sowing these feeds is in March; but if you fow some of them in August, soon after they are ripe, upon a dry soil and in a warm situation, they will abide through the winter, and slower strong the succeeding year; by sowing of the seeds at different times, they may be continued in beauty most parts of the summer." Miller's Gard. Dist. ed. 6. 410.

·Lateral British Covers Agreed Dividing Contract of the second mental enter a man en en de monte de monte de la companya de la co principle and has allow pure they grant to de A Section Line of the ments sond of real sons, or one in the sons face of the sons the west franches and the article of the first in the first of the proad fluor court end when the parties been a provided by donned Milere clay grow the cloth, leaving the three or da lliv spår andrend array a se box lot gri and rower fire this fuccording your by how. begins at the date different times, they see , or continued (sind parts of the fromen," Moso's lord, Diff.





Publish'd by Wartis Botanic Garden Lambeth March.

I Somety del et feute

### TROPÆOLUM MAJUS. GREATER ÎNDIAN-CRESS, OF NASTURTIUM.

\*\*\*\*\*\*

Class and Order.

OCTANDRIA MONOGYNIA.

Generic Character.

Calyx 1-phyllus, calcaratus. Petala 5 in æqualia. Baccæ tres, ficcæ.

Specific Character and Synonyms.

TROPÆOLUM majus foliis peltatis subquinquelobis, petalis obtusis. Lin. Syst. Vegetab. ed. 14.

Murr. p. 357. Sp. Pl. p. 490.


CARDAMINDUM ampliori folio et majori flore. Grande Capucine Tournef. Inst. p. 430.


The present plant is a native of Peru, and is said by Linnæus to have been first brought into Europe in the year 1684; it is certainly one of the greatest ornaments the Flower-Garden can boast: it varies in colour, and is also found in the Nurseries with double slowers. The former, as is well known, is propagated by seed; the latter by cuttings, which should be struck on a hot-bed. To have these plants early, they should be raised with other tender annuals; they usually begin to flower in July, and continue blossoming till the approach of winter: the stalks require to be supported, for if lest to themselves they trail on the ground, overspread, and destroy the neighbouring plants.

Elizabeth Christina, one of the daughters of Linnæus, is faid to have perceived the flowers to emit spontaneously, at certain intervals, sparks like those of electricity, visible only in the dusk of the evening, and which ceased when total darkness came on.

The flowers have the taste of water-cress, with a degree of sweetness, which that plant does not possess, more particularly resident in the spur of the calyx or nectary; hence are sometimes used in fallads, and hence the plant acquires its name of Nasturium.

· Margy is Sign Cath Sergia Correct, or Land Science and ther storic track this prices imprime this prithe the state of t ARDAMIYAMA sushion talle at asjor fare. Charles Louist. Joh. p. 190. retainly one cladic growth ornanciaring it own the can solve a first in three olds of the graphs of relay in the and doubt allow as an entered being a secure to took in a wher ped. The flave their plants couly, the thould be rated with place it clos margin, they ultimally he is flower a July , and common bloffording till the applications wither; document the property of the said the or the second while digits of the evening, and what a control water need carried The flowers have the taffe of we again. The deferred in secured, which that plant short is a pollule, must particulathe related in one fair of the case on resigny should be





I Sonaty Ad a jach.

Publish d by W. Gartis, Botanic Garden, Lambeth March

21

## AGROSTEMMA CORONARIA. ROSE COCKLE, or CAMPION.

\*\*\*\*\*

Class and Order.

DECANDRIA PENTAGYNIA.

Generic Character.

Calyx 1-phyllus, coriaceus. Petala 5 unguiculata: limbo obtufo, indiviso. Caps. 1-locularis.

Specific Character and Synonyms.


AGROSTEMMA coronaria tomentofa, foliis ovato-lanceolatis, petalis emarginatis coronatis ferratis. Lin. Syst. Vegetab. ed. 14. Murr. p. 435. Sp. Pl. p.

LYCHNIS coronaria dioscoridis sativa. Baub. Pin. 203. The single red Rose Campion. Parkins. Parad. p. 252.

Grows spontaneously in Italy and Siberia; Linnæus informs us that the bloffom is naturally white, with red in the middle. " The fingle Rose Campion has been long an inhabitant of " the English gardens, where, by its feeds having scattered, " it is become a kind of weed. There are three varieties of " this plant, one with deep red, another with flesh-coloured, " and a third with white flowers, but these are of small esteem, " for the double Rose Campion being a finer flower, has turned " the others out of most fine gardens. The fingle forts pro-" pagate fast enough by the feeds, the fort with double flowers " never produces any, fo is only propagated by parting of the " roots; the best time for this is in autumn, after their flowers " are past; in doing of this, every head which can be slipped off " with roots should be parted; these should be planted in a " border of fresh undunged earth, at the distance of six inches, " observing to water them gently until they have taken root, " after which they will require no more, for much wet is inju-" rious to them, as is also dung. After the heads are well " rooted, they should be planted into the borders of the " Flower-Garden, where they will be very ornamental during " the times of their flowering, which is in July and August." Miller's Gard. Dict. ed. 6. 4to.

Miller, by mistake, calls this plant Calirofa.

EMERA COO MARIA. ROSE Contract to Ministrac as no and as decreased the sale of the Marginia and a talent de est desire de la comparte del la comparte de la comparte del la comparte de la comparte del la comparte de la compar Commence of the control of the contr





Published by W Curtis Botanic Garden Lamb och March

I Somety del a fat

# DIANTHUS CHINENSIS. CHINA OF INDIAN PINK.

\*\*\*\*\*\*\*\*

Class and Order.
DECANDRIA DIGYNIA.

Generic Character.

Calyx cylindricus, 1-phyllus: basi squamis 4. Petala 5, unguiculata. Capsula cylindrica, 1-locularis.

Specific Character and Synonyms.

DIANTHUS chinensis floribus solitariis, squamis calycinis subulatis patulis, tubum æquantibus, corollis crenatis. Lin. Syst. Vegetab. p. 418. Sp. Pl. 588.

CARYOPHYLLUS finensis supinus, leucoji folio, slore unico. Tournef. act. 1705. p. 348. f. 5.

This species, unknown to the older botanists, is a native of China, hence its name of China Pink; but, in the nurseries, it is in general better known by the name of Indian Pink.


Though it cannot boast the agreeable scent of many of its congeners, it eclipses most of them in the brilliancy of its colours; there are sew slowers indeed which can boast that richness and variety found among the most improved varieties of this species; and as these are easily obtained from seed, so they are found in most collections, both single and double.

It is little better than an annual, but will fometimes continue two years in a dry foil, which it affects.

Attempts have been made to force it, but, as far as we have earned, with no great fuccess.

BARTHES CHITCHES OF CHICA OF Peat. - FOR 150 Charles Charles a meteodo de de la como de la com conserve States followed, formed calcimic relativities or east until firm another in all wel payed and a second and a second and the second and a second and 





I Sowarty; dela faulp

Rublish'd by Warris Betavic Garden Lamb ah Marsh

# STAPELIA VARIEGATA. VARIEGATED STAPELIA.

\*\*\*\*\*\*\*

Class and Order.

PENTANDRIA DIGYNIA.

Generic Character.

Contorta. Nectarium duplici stellula tegente genitalia,

Specific Character and Synonyms.

STAPELIA variegata denticulis ramorum patentibus. Lin. Syst. Vegetab. p. 260. Sp. Pl. p. 316.

ASCLEPIAS aizoides africana. Bradl. fuc. 3. p. 3. t. 22.

This very fingular plant is a native of the Cape of Good Hope, where it grows and flourishes on the rocks with the

Stapelia birfuta.

If these plants be kept in a very moderate stove in winter, and in summer placed in an airy glass-case where they may enjoy much free air, but screened from wet and cold, they will thrive and slower very well; for although they will live in the open air in summer, and may be kept through the winter in a good green-house; yet these plants will not slower so well as those managed in the other way. They must have little water given them, especially in winter.

It is very feldom that the variegata produces feed-veffels in this country; Miller observes, in upwards of forty years that he cultivated it, he never faw it produce its pods but three times, and then on such plants only as were plunged into the


tan-bed in the stove.

This plant may be propagated without feeds, as it grows fast enough from slips; treatment the same as that of the

Creeping Cereus, which fee.

It takes its name of Stapelia from Stapel, a Dutchman, author of some botanical works, particularly a Description of Theophrastus's plants.

T 36 ] its variable ta. Valar of the Bare of the order of the control of property and a second residence of the second second residence of the second 





S. Somerby del. of feutp.

Published by W. Cartis, Botanic Garden Lambeth Marsh.

## CONVOLVULS TRICOLOR. SMALL CONVOL-VULUS OF BINDWEED.

\*\*\*\*\*\*

Class and Order.

PENTANDRIA MONOGYNIA.

Generic Character.

Corolla campanulata, plicata. Stigmata 2. Capfula 2-locu-laris: loculis dispermis.

Specific Character and Synonyms.

CONVOLVULUS tricolor foliis lanceolato ovatis glabris, caule declinato, floribus folitariis. Lin. Syst. Vegetab. 7, 203. Sp. Pl. p. 225.

CONVOLVULUS peregrinus cæruleus, folio oblongo. Baub.

Pin. 295. Flore triplici colore infignito.

Morif. bift. 2. p. 17. f. 1. t. 4. f. 4.


The Spanish Small Blew Bindeweede. Parkins. Parad. p. 4.

This species has usually been called Convolvulus minor by gardeners, by way of distinguishing it from the Convolvulus purpureus, to which they have given the name of major. It is a very pretty annual; a native of Spain, Portugal, and Sicily, and very commonly cultivated in gardens.

The most usual colours of its blossoms are blue, white, and yellow, whence its name of tricolor; but there is a variety of it with white, and another with striped blossoms.

The whole plant with us is in general hairy, hence it does not well accord with Linnæus's description. It is propagated by seeds, which should be sown on the slower-borders in the spring, where the plants are to remain: they require no other care than to be thined and weeded.

Carrier vers Telegrapus Caster The state of the s A CONTRACTOR AND AND A PORTER Applications of the first transport of the supplier of the sup 





Inthinks as the Act directs by M. Ouris Botonic Garden Lombeth Minch 2006.

## PASSIFLORA CŒRULEA. COMMON PASSION-FLOWER.

\*\*\*\*\*\*\*

Class and Order.

GYNANDRIA HEXANDRIA.

Generic Character.

Trigyna. Cal. 5-phyllus. Petala 5. Nectarium corona. Bacca pedicellata.


Specific Character and Synonyms.

PASSIFLORA carulea foliis palmatis integerrimis. Lin. Svst. Vegetab. p. 823. Sp. Pl. p. 1360.

GRANADILLA polyphyllos, fructu ovato. Tourn. inft. 241, FLOS PASSIONIS major pentaphyllus. Sloan. Jam. 104. bift. 1. p. 229.

The Passion-Flower first introduced into this country was the incarnata of Linnæus, a native of Virginia, and figured by Parkinson in his Paradisus Terrestris, who there styles it the surpassing delight of all flowers: the present species, which, from its great beauty and superior hardiness, is now by far the most common, is of more modern introduction; and, though a native of the Brasils, seldom suffers from the severity of our climate; slowering plentifully during most of the summer months, if trained to a wall with a southern aspect, and, in such situations, frequently producing ripe fruit, of the size and form of a large olive, of a pale orange colour.

This most elegant plant may be propagated by seeds, layers, or cuttings; foreign seeds are most to be depended on; they are to be sown in the spring, on a moderate hot-bed, and when the plants are grown to the height of two or three inches, they are to be carefully taken up, and each planted in a separate small pot, silled with good loam, then plunged into a moderate hot-bed, to forward their taking new root; after which they should be gradually inured to the common air: the younger the plants the more shelter they require, and if ever so old or strong, they are in danger from severe frosts. The layers and cuttings are to be treated in the common way, but seedling plants, if they can be obtained, are on many accounts to be preferred.





J. Sowerhy dol. at fault.

## RESEDA ODORATA. SWEET-SCENTED RESEDA OF MIGNONETTE.

\*\*\*\*\*\*\*

Class and Order.

DODECANDRIA TRIGYNIA.

Generic Character.

Cal. 1-phyllus, partitus. Petala laciniata. Caps. ore dehis-

Specific Character and Synonyms.

RESEDA odorata foliis integris trilobisque, calycibus slorem æquantibus. Lin. Syst. Vegetab. p. 449.


RESEDA foliis integris trilobisque, floribus tetragynis.

Mill. Diet. 1. 217.

Mignonette grows naturally in Egypt, it was unknown to the older Botanists; Miller says he received the seeds of it from Dr. Adrian Van Royen, Professor of Botany at Leyden, so that it is rather a modern inhabitant of our gardens.

The luxury of the pleasure-garden is greatly heightened by the delightful odour which this plant diffuses; and as it is most readily cultivated in pots, its fragrance may be conveyed to the parlour of the recluse, or the chamber of the valetudinarian; its persume, though not so refreshing perhaps as that of the Sweet-Briar, is not apt to offend on continuance the most delicate olfactories.

Being an annual it requires to be raifed yearly from feed; when once introduced on a warm dry border it will continue to fow itself, and grow very luxuriantly, flowering from June to the commencement of winter; but as it is defirable to have it as early as possible in the spring, the best way is either to sow the seed in pots in autumn, securing them through the winter in frames, or in a greenhouse, or to raise the seeds early on a gentle hot bed, thinning the plants if they require it, so as to have only two or three in a pot.







Published by W. Ouris Petaric Garden Lambah March

# LILIUM CHALCEDONICUM. CHALCEDONIAN LILY.

\*\*\*\*\*\*\*

### Class and Order.

#### HEXANDRIA MONOGYNIA.

#### Generic Character,

Cor. 6-petala, campanulata: linea longitudinali nectarifera. Capf. valvulis pilo cancellato connexis.

#### Specific Character and Synonyms.

LILIUM chalcedonicum foliis sparsis lanceolatis, sloribus reflexis, corollis revolutis. Lin. Syst. Vegetab. p. 324.

LILIUM byzantium miniatum. Baub. Pin. 78.

The Red Martagon of Constantinople. Park. Parad. p. 34.

This species is best known in the nurseries by the name of the Scarlet Martagon; but as it is not the Martagon of Linnæus, to avoid confusion it will be most proper to adhere to the name which Linnæus has given it.

It is a native not only of Persia, but of Hungary; Prosesfor Jacquin, who has figured it in his most excellent Flora Austriaca, describes it as growing betwixt Carniola and Carinthia, and other parts of Hungary, but always on the tops of the largest mountains.

It varies in the number of its flowers, from one to fix, and

the colour in some is found of a blood red.


Authors differ in their ideas of its smell: Jacquin describing it as disagreeble, while Scopoli compares it to that of an orange.

It flowers in June and July; and is propagated by offsets, which it produces pretty freely, and which will grow in almost

any foil or fituation.

The best time for removing the roots is soon after the leaves are decayed, before they have begun to shoot.

CONTRACTOR OF THE CONTRACTOR 





J. Soverby del Afraile.

Publish d by W. Cartis, Botanic Garden Lambeth Marsh 1707.

# JASMINUM OFFICINALE. COMMON JASMINE or JESSAMINE.

\*\*\*\*\*\*\*

Class and Order.

DIANDRIA MONOGYNIA.

Generic Character.

Cor. 5-fida. Bacca dicocca. Sem. arillata. Antheræ intra tubum,

Specific Character and Synonyms.

JASMINUM officinale foliis oppositis; foliolis distinctis. Lin. Syst. Vegetab. p. 56.

JASMINUM vulgatius flore albo. Baub. Pin. 397. Jasmine or Gesmine. Park. Parad. p. 406.


There is an elegance in the Jasmine which added to its fragrance renders it an object of universal admiration.

"It grows naturally at Malabar, and in feveral parts of India, yet has been long inured to our climate, fo as to thrive and flower extremely well, but never produces any fruit in England. It is easily propagated by laying down the branches, which will take root in one year, and may then be cut from the old plant, and planted where they are designed to remain: it may also be propagated by cuttings, which should be planted early in the autumn, and guarded against the effects of severe frosts.

"When these plants are removed, they should be planted either against some wall, pale, or other sence, where the slexible branches may be supported. These plants should be permitted to grow rude in the summer, otherwise there will be no slowers; but after the summer is past, the luxuriant shoots should be pruned off, and the others must be

" nailed to the fupport.

"There are two varieties of this with variegated leaves, one with white, the other with yellow stripes, but the latter is the most common: these are propagated by budding them on the plain Jasmine; they require to be planted in a warm situation, especially the white-striped, for they are much more tender than the plain, and in very severe winters their branches should be covered with mats or straw to prevent their being killed." Miller's Gard. Dist.





T. MERORITICE SOCIETY OF STREET WELL

and the case of the try but its blottome hally execute in the

## MESEMBRYANTHEMUM DOLABRIFORME. HATCHET-LEAV'D FIG-MARIGOLD.

\*\*\*\*\*\*\*

Class and Order.

ICOSANDRIA PENTAGYNIA.

Generic Character.

Cal. 5-fidus. Petala numerofa, linearia. Capf. carnofa infera, polysperma.

Specific Charaster and Synonyms.

MESEMBRYANTHEMUM dolabriforme acaule, foliis dolabriformibus punctatis. Lin. Syst. Veg. p. 470.

FICOIDES capensis humilis, foliis cornua cervi referentibus, petalis luteis noctiflora, Bradl. suc. 1. p. 11.
t. 10. Dillen Hort. Elth. t. 191. f. 237.


Though many Latin names of plants, as Geranium, Hepatica, Convolvulus, &c. are more familiar to the ear, and more generally used than their English ones, yet Mesembryanthemum though used by some, appears too long to be generally adopted, its English name of Fig-marigold is doubtless to be preferred.

The Fig-marigolds are a very numerous tribe, chiefly inhabitants of the Cape of Good Hope; no less than thirty-three species are figured in that inestimable work the Hortus Eltbamensis of Dillenius. As most of these plants grow readily from slips, or cuttings, and require only the shelter of a common greenhouse, and as they recommend themselves to our notice, either from the extreme singularity of their foliage, the beauty of their slowers, or the peculiarity of their expansion, so they are a savourite class of plants with many.


The present species is a native of the Cape of Good Hope, and is particularly distinguished by having leaves somewhat resembling a hatchet, whence its name; it is as hardy as most, and slowers as freely, but its blossoms fully expand in the evening and night only.

It is very readily propagated by cuttings.

ARROLLER LOOK VINCENSEL WAY . TO 







I worth del a fools

Published as the Act directs by Warris Rotanic Garden Lambeth March.

## ASTER TENELLUS. BRISTLY-LEAV'D ASTER.

\*\*\*\*\*\*

Class and Order.

SYNGENESIA POLYGAMIA SUPERFLUA.

Generic Character.

Recept. nudum. Pappus fimplex. Cor. radii plures 10.

Specific Character and Synonyms. -

- ASTER tenellus foliis subfiliformibus aculeato-ciliatis, pedunculis nudis, calycibus hemisphæricis æqualibus. Lin. Syst. Vegetab. p. 760.
- ASTER parvus æthiopicus, chamæmeli floribus, tamarisci ægyptiaci foliis tenuissime denticulatis. Pluk. alm. 56. t. 271. f. 4. Raii. Suppl. 164. n. 84.


Most of the numerous species of this genus slower about Michaelmas, hence their vulgar name of Michaelmas-Daisy; a name exceptionable not only on account of its length, but from its being a compound word. After, though a Latin term, is now so generally received, that we shall make no apology for adopting it.


We are indebted to North-America for most of our Asters, but the present species, which is omitted by Miller, and is rather a scarce plant in this country, though not of modern introduction, being sigured by Plukenet and described by Ray, is a native of Africa, and, like a few others, requires in the winter the shelter of a greenhouse.


It is particularly diffinguished by having very narrow leaves with short bristles on them, and by its blossoms drooping before they open.

It is a perennial, flowers in September and October, and may be propagated by flips or cuttings.

The plant from whence our drawing was made, came from Messrs. Gordon and Thompson's Nursery, Mile-End.







I Sowerby del. et, faulp.

Publishil by M. Liurte, Retanic Carden Lamboth Morsh.

### [ 34 ]

### BROWALLIA ELATA. TALL BROWALLIA.

\*\*\*\*\*\*\*

Class and Order.

DIDYNAMIA GYMNOSPERMIA.

Generic Character.

Cal. 5-dentatus. Cor. limbus 5-fidus, æqualis, patens: umbilico clauso Antheris 2, majoribus. Caps. 1-locularis.

Specific Character and Synonyms.


BROWALLIA elata pedunculis unifloris multiflorisque. Lin. Syst. Vegetab. p. 572. Sp. Pl. 880. Mill. Dist.

Of this genus there are only two species, both natives of South-America, the elata, so called from its being a much taller plant than the demissa, is a very beautiful, and not uncommon stove or green-house plant; it is impossible, by any colours we have, to do justice to the brilliancy of its slowers.

Being an annual, it requires to be raifed yearly from feed, which must be sown on a hot-bed in the spring, and the plants brought forward on another, otherwise they will not perfect their seeds in this country. Some of these may be transplanted into the borders of the slower-garden which are warmly situated, where, if the season prove savourable, they will slower and ripen their seeds; but, for security's sake, it will be prudent to keep a few plants in the stove or green-house.

As these plants have not been distinguished by any particular English name, MILLER very properly uses its Latin one; a practice which should as much as possible be adhered to, where a genus is named in honour of a Botanist of eminence.


and the state of the state of the say have a subsequence and the state of the collection of the en state in sech . Sylvada add a 31 2 5 6 6 artill violence ..





## [ 35 ]

## CREPIS BARBATA. BEARDED CREPIS, or PURPLE-EYED SUCCORY-HAWKWEED.



Class and Order.

SYNGENESIA POLYGAMIA ÆQUALIS.

#### Generic Character,

Recept. nudum. Cal. calyculatus squamis deciduis. Pappus plumosus, stipitatus.

#### Specific Character and Synonyms,

CREPIS barbata involucris calyce longioribus: fquamis fetaceis sparsis. Lin. Syst. Vegetab. p. 719.

HIERACIUM proliferum falcatum. Baub. Pin. 128.

HIERACIUM calyce barbato. Col. ecpbr. 2. p. 28. t. 27. f. 1.


HIERACIUM boeticum medio nigro. Herm. Parad. Bat. 185. t. 185.

Grows spontaneously in the south of France, about Montpelier; also, in Spain, Italy, Sicily, and elsewhere in the south of Europe: is one of the most common annuals cultivated in our gardens. It begins slowering in July, and continues to blossom till the frost sets in.

No other care is necessary in the cultivation of this species than sowing the seeds in the spring, in little patches, on the borders where they are to remain, thinning them if they prove too numerous.

MILLER calls this species bætica, and improperly describes the centre of the flower as black, as also does HERMAN: in all the specimens we have seen, it has evidently been of a deep purple colour, or, as LINNEUS expresses it, atropurpurascens.

Least to the state of the Stanfaller . Manager of the same of the sam 





Harriside et faulp.

Published by M. On the Belanie Ger den Lambate March

## LILIUM BULBIFERUM. ORANGE LILY.

\*\*\*\*\*\*

Class and Order.

#### HEXANDRIA MONOGYNIA.

Generic Character.

Cor. 6-petala, campanulata: linea longitudinali nectarifera. Capf. valvulis pilo cancellato connexis.

Specific Character and Synonyms.

LILIUM bulbiferum foliis sparsis, corollis campanulatis erectis: intus scabris. Lin. Syst. Vegetab. p. 324. Jacq. Fl. Austr. t. 226.

LILIUM purpureo-croceum majus. Baub. Pin. 76. LILIUM aureum, the gold red Lily. Park. Parad. p. 37.

" The fmaller orange Lily.

Bears the smoke of London better than many plants. Varies with and without bulbs on the stalks.

<sup>&</sup>quot;The common orange or red Lily is as well known in the English gardens as the white Lily, and has been as long cultivated here. This grows naturally in Austria and some parts of Italy. It multiplies very fast by offsets from the roots, and is now so common as almost to be rejected; however, in large gardens these should not be wanting, for they make a good appearance when in flower, if they are properly disposed; of this fort there are the following varieties:

<sup>&</sup>quot;The orange Lily with double flowers,
"The orange Lily with variegated leaves,

<sup>&</sup>quot;These varieties have been obtained by culture, and are preserved in the gardens of florists. They all flower in June and July, and their stalks decay in September, when the roots may be transplanted and their offsets taken off, which should be done once in two or three years, otherwise their branches will be too large, and the flower-stalks weak. This doth not put out new roots till towards spring, fo that the roots may be transplanted any time after the stalks decay till November. It will thrive in any soil or situation, but will be strongest in a soft gentle loam, not too moist." Mill. Diet.

JE INSTITUTE HOLDE -----And the second states of the second s principle to the state of the second of the of the state of th